GREEN•SOLTM SUL-15 PLUS

ACTIVE INGREDIENT:

Gibberellic Acid

GUARANTEED ANALYSIS (Fertilizer)

Sulfur (combined) as S ------15.00%

Derived from Liquid Ammonium Thiosulfate.

GENERAL INFORMATION

Sulfur is a chemical element and as a fertilizer has many essential roles in plants. SUL-15 PLUS LIQUID activates the enzyme and hormone systems in plants which are essential for normal plant growth and reproduction. SUL-15 PLUS LIQUID provides nutrients for healthier roots; stronger stems; stronger, longer stalks; better fruit; and improved plant performance. SUL-15 PLUS LIQUID is beneficial when applied anytime during the growing season.

METHOD OF APPLICATION

Unless otherwise specified, dosage rates are to be mixed in 5 to 10 gallons of water as a foliar spray. Spray at 25-40 lbs. of pressure with No. 1.5 or larger flood tips, or other tips that provide a fine spray for complete coverage. Application at temperatures above 85°F and with low humidity should have a no-drift product added to insure proper coverage. SUL-15 PLUS LIQUID should always be the last ingredient added to tank mix.

COMPATIBILITY

SUL-15 PLUS LIQUID is compatible with pesticides used commonly in vegetable and fruit crop production.

ATTENTION

Do not use with dinitro compounds, dormant oil, Bordeaux, spray lime or highly alkaline spray materials. Always add to the spray mixture as the last ingredient.

APPLICATION WITH FERTILIZER

SUL-15 PLUS LIQUID may be added with 32% UAN solution or may be impregnated on granular fertilizer at a rate of 1 quart per 300 lbs.

SUGGESTED RATES

COTTON: One quart/acre broadcast or 1 pint/acre banded directed to foliage at 4-6 leaf stage. Broadcast 1 quart/acre at or prior to pin head square. Apply 1 pint/acre with each application of foliar applied urea and insecticides.

SOYBEANS: Broadcast 1 quart/acre at 3rd trifoliate leaf stage. Broadcast 1 quart/acre prior to blooming. Apply 1 pint/acre to insecticide sprays to promote increased pod fill.

PEANUTS: Band 1 quart/acre at first true leaf. Apply 1 pint/acre with each application of fungicide.

CEREAL OR SMALL GRAINS (WHEAT, OATS, BARLEY, RICE, RYE): Apply 1 quart/acre with nitrogen application.

COARSE GRAINS (CORN, GRAIN SORGHUM): Broadcast 1 quart/acre at 3-4 leaf stage. Broadcast 1 quart/acre at 7-8 leaf stage.

BEANS: Broadcast 1 quart/acre at 3rd trifoliate leaf stage. Broadcast 1 quart/acre prior to blooming. Apply 1 pint/acre with insecticide sprays. CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS: Apply 1 quart/acre at 2-3" stage. Repeat at 14-day intervals during the growing season.

SQUASH, CUCUMBERS: Apply 1 quart/acre at bunching. Repeat at 14-day intervals during the growing season. LEAFY VEGETABLES (LETTUCE, SPINACH, GREENS): Apply 1 quart/acre at 2-3 inch stage. Repeat at 14-day intervals during the growing season. For best results use a non-toxic surfactant or food grade spreader sticker.

MELONS (WATERMELONS, CANTALOUPES, ETC.): Apply 1 quart/acre at bunching. Repeat at 14-day intervals during the growing season. STRAWBERRIES, TOMATOES, PEPPERS: Apply 1 quart/acre 7 days after planting. Repeat at 14-day intervals during the growing season. POTATOES: Apply 1 quart/acre at the 2-4 leaf stage. Repeat at 14-day intervals during the growing season.

SUGARBEETS: Apply 1 quart/acre at 2-4 leaf stage. Repeat application in 3-4 weeks. Make final application of 2 quarts/acre 3-4 weeks later. SUGARCANE: Apply 1 quart/acre 6-8 weeks after planting or harvest. Reapply every 6-8 weeks during the growing season.

CITRUS (ORANGES, LEMONS, GRAPEFRUIT): Apply 2-4 quarts in 100 gallons of water at 14-day intervals during the growing season. Cease applications 2 weeks prior to harvest.

OUTDOOR/INDOOR FLOWERING SHRUBS, ORNAMENTAL PLANTS: Apply 2 oz. per gallon of water sprayed in a fine mist every 14 days during the growing season. (DO NOT USE ON FERNS.)

MANUFACTURED BY: FRIT INDUSTRIES JODIE PARKER ROAD **OZARK, AL 36360**

EPA EST. NO. 64872-AL-001 EPA REG. NO. 64872-1 NET WEIGHT: 2.5 GAL (24.25 LBS.)