

# GREEN•SOL™

## SUL-15 PLUS POWDER

SULFUR (Combined) as S -----	15.00%
BORON (B) -----	0.02%
IRON (Fe)-----	0.10%
0.10% Chelated Iron	

**Derived From: Ammonium Sulfate, Iron Chelate, And Sodium Borate. Chelating Agent Is Ethylenediamine tetra-acetate.**

### GENERAL INFORMATION

Sulfur is an essential element and as such has many essential roles in plants. SUL-15 PLUS POWDER activates certain enzyme and hormone systems in plants which are essential for normal plant growth and reproduction. SUL-15 PLUS POWDER provides nutrients for healthier roots, stronger stems and stalks and better fruit. SUL-15 PLUS POWDER can be applied anytime during the growing season.

### METHOD OF APPLICATION

Unless otherwise specified, the product is to be mixed in 5 to 10 gallons of water and applied as a foliar spray. Spray at 25-40 lbs. of pressure with No. 1.5 or larger flood tips, or other tips that provide a fine spray for complete coverage. Application at temperatures above 85°F and with low humidity should have a no-drift product added to insure proper coverage. SUL-15 PLUS POWDER should always be the last ingredient added to tank mix.

### COMPATIBILITY

SUL-15 PLUS POWDER is compatible with pesticides commonly used in vegetable and fruit crop production.

### ATTENTION

Do not use with dinitro compounds, dormant oil, Bordeaux, spray lime or highly alkaline spray materials. Always add to the spray mixture as the last ingredient.

**APPLICATION WITH FERTILIZER:** SUL-15 PLUS POWDER may be applied with 32% UAN solution.

### SUGGESTED RATES

**COTTON:** Broadcast 2#/acre to foliage at 4-6 leaf stage. Broadcast 2#/acre at or prior to pin head square. Apply 1#/acre with each application of foliar applied urea and insecticides.

**SOYBEANS:** Broadcast 2#/acre at 3rd trifoliate leaf stage. Broadcast 2#/acre prior to blooming. Apply 1 #/acre with insecticide sprays to promote increased pod fill.

**PEANUTS:** Band 2#/acre at first true leaf. Apply 1 #/acre with each application of fungicide (5-7 times).

**CEREAL OR SMALL GRAINS (WHEAT, OATS, BARLEY, RICE, RYE):** Apply 2#/acre with nitrogen application.

**COARSE GRAINS (CORN, GRAIN SORGHUM):** Broadcast 2#/acre at 3-4 leaf stage. Broadcast 2#/acre at 7-8 leaf stage.

**CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS:** Apply 2#/acre at 2-3" stage. Repeat at 14-day intervals during the growing season.

**SQUASH, CUCUMBERS:** Apply 2#/acre at bunching. Repeat at 14-day intervals during the growing season.

**LEAFY VEGETABLES (LETTUCE, SPINACH, GREENS):** Apply 2#/acre at 2-3 inch stage. Repeat at 14-day intervals during the growing season. For best results use a non-toxic surfactant or food grade spreader sticker.

**MELONS (WATERMELONS, CANTALOUPE, ETC.):** Apply 2#/acre at bunching. Repeat at 14-day intervals during the growing season.

**STRAWBERRIES, TOMATOES, PEPPERS:** Apply 2#/acre 7 days after planting. Repeat at 14-day intervals during the growing season.

**POTATOES:** Apply 2#/acre at the 2-4 leaf stage. Repeat at 14-day intervals during the growing season.

**SUGARBEETS:** Apply 2#/acre at 2-4 leaf stage. Repeat application in 3-4 weeks. Make final application of 4#/acre 3-4 weeks later.

**SUGARCANE:** Apply 2#/acre 7-14 days after planting.

**CITRUS (ORANGES, LEMONS, GRAPEFRUIT):** Apply 4-8 pounds in 100 gallons of water at 14-day intervals during the growing season. Cease applications 2 weeks prior to harvest.

**OUTDOOR/INDOOR FLOWERING SHRUBS, ORNAMENTAL PLANTS:** Apply 4 Tablespoons per gallon of water sprayed in a fine mist every 14 days during the growing season. **(DO NOT USE ON FERNS.)**

**NET WEIGHT: 25 LBS. (11.35 KG)**

**MANUFACTURED FOR:**  
**GREEN•SOL INCORPORATED**  
**PO BOX 1589, OZARK, AL 36361-1589**